antibodies .- online.com







FLT3 Protein (AA 27-541) (Fc Tag)



Images



Overview


Quantity:	100 μg
Target:	FLT3
Protein Characteristics:	AA 27-541
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This FLT3 protein is labelled with Fc Tag.

Product Details

Purpose:	Human FLT3/Flk-2 Protein
Sequence:	Asn27-Asn541
Characteristics:	Recombinant Human FLT3/Flk-2 Protein is expressed from HEK293 with hFc tag at the C-Terminus.It contains Asn27-Asn541.
Purity:	> 95 % as determined by Tris-Bis PAGE
Sterility:	0.22 μm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	Immobilized Human FLT3 Ligand, His Tag at 1µg/ml (100µl/Well) on the plate. Dose response curve for Human FLT3, hFc Tag with the EC50 of 19.6ng/ml determined by ELISA. See testing image for detail.


Target Details

Target:	FLT3
Alternative Name:	FLT3 (FLT3 Products)
Background:	The Flt-3 (fms-like tyrosine kinase) receptor, also named Flk-2 (fetal liver kinase) and Stk-1 (stem cell tyrosine kinase) is a member of the class III subfamily of receptor tyrosine kinases that also includes KIT, the receptor for SCF and FMS, the receptor for M-CSF. Tyrosine-protein kinase that acts as cell-surface receptor for the cytokine FLT3LG and regulates differentiation, proliferation and survival of hematopoietic progenitor cells and of dendritic cells. Promotes phosphorylation of SHC1 and AKT1, and activation of the downstream effector MTOR. Promotes activation of RAS signaling and phosphorylation of downstream kinases, including MAPK1/ERK2 and/or MAPK3/ERK1.
Molecular Weight:	84.9 kDa. Due to glycosylation, the protein migrates to 100-120 kDa based on Tris-Bis PAGE result.
Pathways:	RTK Signaling
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from $0.22\mu m$ filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months




SDS-PAGE

Image 1. Human FLT3 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 %.



Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Human FLT3/Flk-2 is greater than 95 % as determined by SEC-HPLC.



SDS-PAGE

 $\label{eq:mage_3.} \textbf{Image 3.} \ \ \textbf{Human FLT3/Flk-2} \ \ \textbf{on Tris-Bis PAGE under}$ reduced condition. The purity is greater than 95 % .

Please check the product details page for more images. Overall 5 images are available for ABIN7274657.