antibodies - online.com







FOLR1 Protein (AA 25-233) (His-Avi Tag)





Overview

Quantity:	100 μg
Target:	FOLR1
Protein Characteristics:	AA 25-233
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This FOLR1 protein is labelled with His-Avi Tag.

Product Details


Purpose:	Human FOLR1 Protein
Sequence:	Arg25-Met233
Characteristics:	Recombinant Human FOLR1 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Arg25-Met233.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	Immobilized Human FOLR1, His Tag at 0.5µg/ml (100µl/Well) on plate. Dose response curve for Anti-FOLR1 Antibody, hFc Tag with the EC50 of 15ng/ml determined by ELISA. See testing image for detail.

Target Details

(FBP), is a 37 - 42 kDa protein that mediates the cellular uptake of folic acid and reduced folates. Dietary folates are required for many key metabolic processes including nucleotic methionine synthesis, the interconversion of glycine and serine, and histidine breakdown FOLR1 binds to folate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate and folate analogs into the interior of cells. Has high affinity for for and folic acid analogs at neutral pH . Molecular Weight: 27.5 kDa. Due to glycosylation, the protein migrates to 38-45 kDa based on Tris-Bis PAGE UniProt: P15328 Pathways: Dicarboxylic Acid Transport Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is addiprotectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after	rarget Details	
Background: Folate Receptor 1 (FOLR1), also known as Folate Receptor alpha and Folate Binding Prot (FBP), is a 37 - 42 kDa protein that mediates the cellular uptake of folic acid and reduced folates. Dietary folates are required for many key metabolic processes including nucleoti methionine synthesis, the interconversion of glycine and serine, and histidine breakdown FOLR1 binds to folate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate and folate analogs into the interior of cells. Has high affinity for for and folic acid analogs at neutral pH . Molecular Weight: 27.5 kDa. Due to glycosylation, the protein migrates to 38-45 kDa based on Tris-Bis PAGE UniProt: P15328 Pathways: Dicarboxylic Acid Transport Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is addiprotectant before lyophilization. Storage: -20 °C-80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the proteir smaller quantities for optimal storage. Please minimize freeze thaw cycles.	Target:	FOLR1
(FBP), is a 37 - 42 kDa protein that mediates the cellular uptake of folic acid and reduced foliates. Dietary foliates are required for many key metabolic processes including nucleotic methionine synthesis, the interconversion of glycine and serine, and histidine breakdown FOLR1 binds to foliate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate and foliate analogs into the interior of cells. Has high affinity for for and folic acid analogs at neutral pH . Molecular Weight: 27.5 kDa. Due to glycosylation, the protein migrates to 38-45 kDa based on Tris-Bis PAGE UniProt: P15328 Pathways: Dicarboxylic Acid Transport Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening, Reconstituting to a concentration more than 100 µg/recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is addiprotectant before lyophilization. Storage: -20 °C,80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the proteir smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	Alternative Name:	FOLR1 (FOLR1 Products)
UniProt: P15328 Pathways: Dicarboxylic Acid Transport Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	Background:	folates. Dietary folates are required for many key metabolic processes including nucleotide and methionine synthesis, the interconversion of glycine and serine, and histidine breakdown. FOLR1 binds to folate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate and folate analogs into the interior of cells. Has high affinity for folate
Pathways: Dicarboxylic Acid Transport Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C, 80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	Molecular Weight:	27.5 kDa. Due to glycosylation, the protein migrates to 38-45 kDa based on Tris-Bis PAGE result
Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt., -80 °C for 3-6 months after reconstitution., 2-8 °C for 2-7 days after reconstitution., Recommend to aliquot the protein smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	UniProt:	P15328
Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt., -80 °C for 3-6 months after reconstitution., 2-8 °C for 2-7 days after reconstitution., Recommend to aliquot the protein smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	Pathways:	Dicarboxylic Acid Transport
Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt., -80 °C for 3-6 months after reconstitution., 2-8 °C for 2-7 days after reconstitution., Recommend to aliquot the protein smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	Application Details	
Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	Restrictions:	For Research Use only
Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	Handling	
recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	Format:	Lyophilized
protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt., -80 °C for 3-6 months after reconstitution., 2-8 °C for 2-7 days after reconstitution., Recommend to aliquot the protein smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	Buffer:	Lyophilized from $0.22\mu m$ filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	Storage:	-20 °C,-80 °C
Expiry Date: 12 months	Storage Comment:	reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into
	Expiry Date:	12 months

Human FOLR1, His Tag ELISA

0.05µg Human FOLR1, His Tag Per Well




ELISA

Image 1. Immobilized Human FOLR1, His Tag at $0.5 \,\mu g/mL$ (100 $\,\mu L/Well$) on plate.Dose response curve for Anti-FOLR1 Antibody, hFc Tag with the EC50 of 15 ng/mL determined by ELISA.

Human FOLR1, His Tag ELISA

0.05µg Human FOLR1, His Tag Per Well



ELISA

Image 2. Immobilized Human FOLR1, His Tag at $0.5 \,\mu\text{g/mL}$ (100 $\,\mu\text{L/Well}$) on plate.Dose response curve for Anti-FOLR1 Antibody, hFc Tag with the EC50 of 11.9 ng/mL determined by ELISA.

140KD 115KD 80KD 70KD 50KD 40KD 30KD 25KD

10KD

MK

R

SDS-PAGE

Image 3. Human FOLR1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .

Please check the product details page for more images. Overall 4 images are available for ABIN7274667.