antibodies

Datasheet for ABIN7274673 FCGR1 Protein (AA 16-288) (His-Avi Tag,Biotin)



4 Images



Quantity:	100 µg
Target:	FCGR1
Protein Characteristics:	AA 16-288
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This FCGR1 protein is labelled with His-Avi Tag,Biotin.


Product Details


Purpose:	Biotinylated Human Fc gamma RI/CD64 Protein
Sequence:	GIn16-Pro288
Characteristics:	Recombinant Biotinylated Human Fc gamma RI/CD64 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.It contains GIn16-Pro288.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per μ g by the LAL method.
Biological Activity Comment:	The affinity constant of 0.91 nM as determined in SPR assay (Biacore T200). See testing image for detail.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7274673 | 01/18/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Target:	FCGR1
Alternative Name:	Fc gamma RI (FCGR1 Products)
Background:	The Fc gamma Rs have been divided into three classes based on close relationships in their extracellular domains, these groups are designated Fc gamma RI (also known as CD64), Fc gamma RII (CD32), and Fc gamma RIII (CD16). Each group may be encoded by multiple genes and exist in different isoforms depending on species and cell type.
Molecular Weight:	33.5 kDa. Due to glycosylation, the protein migrates to 55-68 kDa based on Tris-Bis PAGE result
Pathways:	Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months






Surface Plasmon Resonance

Image 1. Biotinylated Human Fc gamma RI, His-Avi Tag captured on CM5 Chip via Anti-His Antibody can bind Trastuzumab with an affinity constant of 0.91 nM as determined in SPR assay (Biacore T200).

SDS-PAGE

Image 2. Biotinylated Human Fc gamma RI/CD64 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .





ELISA

Image 3. Immobilized Anti-HER2 Antibody, hFc Tag at 0.5μ g/mL (100 μ L/Well) on the plate. Dose response curve for Biotinylated Human Fc gamma RI, His Tag with the EC50 of 4.7 ng/mL determined by ELISA.

Log Biotinylated Human Fc gamma RI, His Tag Conc.(µg/ml)

Please check the product details page for more images. Overall 4 images are available for ABIN7274673.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7274673 | 01/18/2024 | Copyright antibodies-online. All rights reserved.