antibodies - online.com







HLAG Protein (Monomer) (HLA-G)

Images



Overview

Quantity:	100 μg
Target:	HLAG
Protein Characteristics:	Monomer
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This HLAG protein is labelled with HLA-G.

Product Details


Purpose:	Cynomolgus HLA-G & B2M & Peptide (RIIPRHLQL) Monomer Protein
Sequence:	Gly25-Thr305 (HLA-G), Ile21-Met119 (B2M) and RIIPRHLQL peptide
Specificity:	Uni-Prot: E0WKX9 (HLA-G), Q8SPW0 (B2M), RIIPRHLQL
Characteristics:	Recombinant Cynomolgus HLA-G & B2M & Peptide (RIIPRHLQL) Monomer Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Gly25-Thr305(HLA-G), Ile21-Met119(B2M) and RIIPRHLQL peptide.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.

Target Details

Expiry Date:


12 months

Target:	HLAG
Alternative Name:	HLA-G (HLAG Products)
Background:	HLA-G is a molecule that was first known to confer protection to the fetus from destruction by the immune system of its mother, thus critically contributing to fetal-maternal tolerance. The first functional finding constituted the basis for HLA-G research and can be summarized as such: HLA-G, membrane-bound or soluble, strongly binds its inhibitory receptors on immune cells (NK, T, B, monocytes/dendritic cells), inhibits the functions of these effectors, and so induces immune inhibition.
Molecular Weight:	50.6 kDa. Due to glycosylation, the protein migrates to 55-65 kDa based on Tris-Bis PAGE result
UniProt:	E0WKX9
Pathways:	Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Cancer Immune Checkpoints
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after



SDS-PAGE

Image 1. Cynomolgus HLA-G on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .



Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Cynomolgus HLA-G is greater than 95 % as determined by SEC-HPLC.