# antibodies -online.com







### **HLAG Protein (Monomer) (HLA-G)**

**Images** 



#### Overview

Quantity:	100 μg
Target:	HLAG
Protein Characteristics:	Monomer
Origin:	Rhesus Monkey
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This HLAG protein is labelled with HLA-G.

#### **Product Details**

Purpose:	Rhesus macaque HLA-G & B2M & Peptide (RIIPRHLQL) Monomer Protein
Sequence:	Gly25-Thr305 (HLA-G), Ile21-Met119 (B2M) and RIIPRHLQL peptide
Specificity:	Uni-Prot: 002948 (HLA-G), Q6V7J5 (B2M), RIIPRHLQL
Characteristics:	Recombinant Rhesus macaque HLA-G & B2M & Peptide (RIIPRHLQL) Monomer Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Gly25-Thr305(HLA-G), Ile21-Met119(B2M) and RIIPRHLQL peptide.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per μg by the LAL method.

Storage:

Expiry Date:

Storage Comment:

Target Details	
Target:	HLAG
Alternative Name:	HLA-G (HLAG Products)
Background:	HLA-G is a molecule that was first known to confer protection to the fetus from destruction by the immune system of its mother, thus critically contributing to fetal-maternal tolerance. The first functional finding constituted the basis for HLA-G research and can be summarized as such: HLA-G, membrane-bound or soluble, strongly binds its inhibitory receptors on immune cells (NK, T, B, monocytes/dendritic cells), inhibits the functions of these effectors, and so induces immune inhibition.
Molecular Weight:	50.5 kDa. Due to glycosylation, the protein migrates to 58-63 kDa based on Tris-Bis PAGE result.
UniProt:	002948
Pathways:	Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process,  Cancer Immune Checkpoints
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as

-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after


smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into

protectant before lyophilization.


-20 °C,-80 °C

12 months



#### **SDS-PAGE**

Image 1. Rhesus macaque HLA-G on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .



## Size-exclusion chromatography-High Pressure Liquid Chromatography

**Image 2.** The purity of Rhesus macaque HLA-G is greater than 95 % as determined by SEC-HPLC.