

Datasheet for ABIN7274834

**ICAM1 Protein (AA 28-480) (His tag)****2** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	ICAM1
Protein Characteristics:	AA 28-480
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ICAM1 protein is labelled with His tag.

## Product Details

Purpose:	Cynomolgus ICAM-1/CD54 Protein
Sequence:	Gln28-Val480
Characteristics:	Recombinant Cynomolgus ICAM-1/CD54 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Gln28-Val480.
Purity:	> 95 % as determined by Tris-Bis PAGE, > 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.

## Target Details

Target:	ICAM1
Alternative Name:	ICAM-1 ( <a href="#">ICAM1 Products</a> )

## Target Details


Background:	Intercellular adhesion molecule-1 (ICAM-1, CD54) is a 90 kDa member of the immunoglobulin (Ig) superfamily and is critical for the firm arrest and transmigration of leukocytes out of blood vessels and into tissues. ICAM-1 is constitutively present on endothelial cells, but its expression is increased by proinflammatory cytokines. The endothelial expression of ICAM-1 is increased in atherosclerotic and transplant-associated atherosclerotic tissue and in animal models of atherosclerosis.
Molecular Weight:	50.55 kDa. Due to glycosylation, the protein migrates to 75-100 kDa based on Tris-Bis PAGE result.
UniProt:	<a href="#">A0A2K5UKF7</a>
Pathways:	<a href="#">Cellular Response to Molecule of Bacterial Origin</a> , <a href="#">Regulation of Actin Filament Polymerization</a> , <a href="#">Carbohydrate Homeostasis</a> , <a href="#">Regulation of Leukocyte Mediated Immunity</a> , <a href="#">Thromboxane A2 Receptor Signaling</a>

## Application Details

Restrictions:	For Research Use only
---------------	-----------------------


## Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS ( pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months



SDS-PAGE

**Image 1.** Cynomolgus ICAM-1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .



Size-exclusion chromatography-High Pressure Liquid Chromatography

**Image 2.** The purity of Cynomolgus ICAM-1 is greater than 95 % as determined by SEC-HPLC.