.-online.com antibodies

Datasheet for ABIN7274936 IL-31 Protein (AA 27-163) (His tag)

3 Images



Overview


| Quantity: | 100 µg |
|-------------------------------|--|
| Target: | IL-31 (IL31) |
| Protein Characteristics: | AA 27-163 |
| Origin: | Cynomolgus |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This IL-31 protein is labelled with His tag. |

Product Details

| Purpose: | Cynomolgus IL-31 Protein |
|------------------------------|--|
| Sequence: | Leu27-Thr163 |
| Characteristics: | Recombinant Cynomolgus IL-31 Protein is expressed from HEK293 with His tag at the N- Terminus.It contains Leu27-Thr163. |
| Purity: | > 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC |
| Sterility: | 0.22 µm filtered |
| Endotoxin Level: | Less than 1EU per μ g by the LAL method. |
| Biological Activity Comment: | The affinity constant of 0.715 nM as determined in SPR assay (Biacore T200). See testing image for detail. |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7274936 | 01/18/2024 | Copyright antibodies-online. All rights reserved.

| Target Details | |
|---------------------|---|
| Target: | IL-31 (IL31) |
| Alternative Name: | IL-31 (IL31 Products) |
| Background: | Many autoimmune skin diseases, such as bullous pemphigoid (BP), psoriasis and certain types of chronic urticaria, are associated with intensive pruritus. While histamine and neuropeptides have previously been ascribed to play a role in itch that accompanies these diseases, recent evidence suggests that the pruritogenic cytokine interleukin (IL)-31 is a major driver of pruritic responses. IL-31 was originally shown to be produced by activated helper T cells, particularly Th2 cells, mast cells, macrophages and dendritic cells. |
| Molecular Weight: | 16.5 kDa. Due to glycosylation, the protein migrates to 23-30 kDa based on Tris-Bis PAGE result. |
| Application Details | |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Reconstitution: | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water. |
| Buffer: | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |
| Expiry Date: | 12 months |




SDS-PAGE

Image 1. Cynomolgus IL-31 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .

Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Cynomolgus IL-31 is greater than 95 % as determined by SEC-HPLC.



Surface Plasmon Resonance

Image 3. Human IL-31 RA, hFc Tag captured on CM5 Chip via Protein A can bind Cynomolgus IL-31, His Tag with an affinity constant of 0.715 nM as determined in SPR assay (Biacore T200).

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7274936 | 01/18/2024 | Copyright antibodies-online. All rights reserved.