# antibodies .- online.com







## **IL-7 Protein**





#### Overview

Quantity:	100 μg
Target:	IL-7 (IL7)
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant

## **Product Details**

Sequence:	Asp26-His177
Purity:	> 95% as determined by Tris-Bis PAGE,> 95% as determined by HPLC
Sterility:	0.22 μm filtered
Endotoxin Level:	Less than 1EU per μg by the LAL method.
Biological Activity Comment:	Immobilized Human IL-7 at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Human IL-7R alpha, hFc Tag with the EC50 of 0.21µg/ml determined by ELISA. The affinity constant of 22.66 nM as determined in SPR assay (Biacore T200). See testing image for detail.

## **Target Details**

Target:	IL-7 (IL7)
Alternative Name:	IL-7 (IL7 Products)
Background:	IL7, Interleukin-7, IL-7, Lymphopoietin-1, PBGF,Interleukin-7 (IL7) plays a nonredundant role in T
	cell survival and homeostasis, which is illustrated in the severe T cell lymphopenia of IL7- deficient mice, or demonstrated in animals or humans that lack expression of either the IL7Ra

## **Target Details**

	or $\gamma$ c chain, the two subunits that constitute the functional IL7 receptor.
Molecular Weight:	17.5 kDa same as Tris-Bis PAGE result.
UniProt:	P13232
Pathways:	JAK-STAT Signaling

## **Application Details**


|--|--|--|

## Handling

Format:	Lyophilized
Reconstitution:	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 $\mu$ g/mL is recommended (usually we use 1 mg/mL solution for lyophilization). Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	4 °C,-80 °C
Storage Comment:	Reconstituted protein stable at -80°C for 12 months, 4°C for 1 week. Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Expiry Date:	12 months

## **Images**


## Human IL-7, No Tag ELISA 0.05µg Human IL-7, No Tag Per Well



#### **ELISA**

**Image 1.** Immobilized Human IL-7 at  $0.5\,\mu g/mL$  ( $100\,\mu$  L/Well) on the plate. Dose response curve for Human IL-7R alpha, hFc Tag with the EC50 of  $0.21\,\mu g/mL$  determined by ELISA.

#### IL-7 Stimulation PHA-pretreated PBMC proliferation assay



40KD 30KD 25KD

15KD

10KD

#### **Proliferation Assay**

**Image 2.** Human IL-7, No Tag stimulates proliferation of PHA-pretreated human peripheral blood mononuclear cell (PBMC) with the EC50 for this effect is 3.5 ng/mL.

#### **SDS-PAGE**

**Image 3.** Human IL-7 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

Please check the product details page for more images. Overall 6 images are available for ABIN7274963.