

Datasheet for ABIN7274976

IL18BP Protein (AA 31-194) (Fc Tag)**5** Images[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	IL18BP
Protein Characteristics:	AA 31-194
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL18BP protein is labelled with Fc Tag.

Product Details

Purpose:	Human IL-18BP Protein
Sequence:	Thr31-Gly194
Characteristics:	Recombinant Human IL-18BP Protein is expressed from HEK293 with hFc tag at the C-Terminus. It contains Thr31-Gly194.
Purity:	> 95 % as determined by Tris-Bis PAGE
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	Immobilized Human IL-18 at 5µg/ml (100µl/Well) on the plate. Dose response curve for Human IL-18BP, hFc Tag with the EC50 of 18.2ng/ml determined by ELISA. The affinity constant of 0.04nM as determined in SPR assay (Biacore T200). See testing image for detail.

Target Details

Target:	IL18BP
Alternative Name:	IL-18BP (IL18BP Products)
Background:	Cytokines were the first modern immunotherapies to produce durable responses in patients with advanced cancer, components of the interleukin-18 (IL-18) pathway are upregulated on tumour-infiltrating lymphocytes, suggesting that IL-18 therapy could enhance anti-tumour immunity. IL-18BP, a high-affinity IL-18 decoy receptor, is frequently upregulated in diverse human and mouse tumours and limits the anti-tumour activity of IL-18 in mice.
Molecular Weight:	44.4 kDa. Due to glycosylation, the protein migrates to 60-80 kDa based on Tris-Bis PAGE result.



Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C, -80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months

Human IL-18BP, hFc Tag ELISA
0.5µg Human IL-18, No Tag Per Well



ELISA


Image 1. Immobilized Human IL-18 at 5 µg/mL (100 µL/Well) on the plate. Dose response curve for Human IL-18BP, hFc Tag with the EC50 of 18.2 ng/mL determined by ELISA.

SDS-PAGE

Image 2. Human IL-18BP on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .

Surface Plasmon Resonance

Image 3. Human IL-18BP, hFc Tag captured on Protein A chip, can bind Human IL-18 with an affinity constant of 0.04nM as determined in a SPR assay(Biacore T200).



Please check the [product details page](#) for more images. Overall 5 images are available for ABIN7274976.