antibodies - online.com







IL-9 Protein (AA 19-144) (Fc Tag)



Images



Overview


Quantity:	100 μg
Target:	IL-9 (IL9)
Protein Characteristics:	AA 19-144
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL-9 protein is labelled with Fc Tag.

Product Details


Purpose:	Human IL-9 Protein
Sequence:	Gln19-Ile144
Characteristics:	Recombinant Human IL-9 Protein is expressed from HEK293 with hFc tag at the C-Terminus.It contains Gln19-Ile144.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	Immobilized Human IL-9, hFc Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve
	for Biotinylated Anti-IL-9 Antibody, hFc Tag with the EC50 of 14.5ng/ml determined by ELISA.
	See testing image for detail.


Target Details

. a. gat 2 atama	
Target:	IL-9 (IL9)
Alternative Name:	IL-9 (IL9 Products)
Background:	IL-9 is a pleiotropic cytokine that influences various distinct functions of different target cells such as T cells, B cells, mast cells and airway epithelial cells by activating STAT1, STAT3 and STAT5. Because of its pleiotropic functions, IL-9 has been demonstrated to be involved in several diseases, such as cancer, autoimmunity and other pathogen-mediated immune-regulated diseases. In this review, we focus on the role of Th9 and IL-9-producing cells in allergic asthma.
Molecular Weight:	40.9 kDa. Due to glycosylation, the protein migrates to 50-70 kDa based on Tris-Bis PAGE result.
UniProt:	P15248
Pathways:	JAK-STAT Signaling
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from $0.22\mu m$ filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months



Log Biotinylated Anti-IL-9 Antibody, hFc Tag Conc.(µg/ml)





ELISA

Image 1. Immobilized Human IL-9, hFc Tag at $0.5\,\mu\text{g/mL}$ (100 $\mu\text{L/Well}$) on the plate. Dose response curve for Biotinylated Anti-IL-9 Antibody, hFc Tag with the EC50 of 14.5 ng/mL determined by ELISA.

Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Human IL-9 is greater than 95 % as determined by SEC-HPLC.

SDS-PAGE

Image 3. Human IL-9 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .