antibodies -online.com





IL23R Protein (AA 24-355) (mFc Tag)





Go to Product page

Overview

Quantity:	100 μg
Target:	IL23R
Protein Characteristics:	AA 24-355
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL23R protein is labelled with mFc Tag.

Product Details


Purpose:	Human IL-23R Protein
Sequence:	Gly24-Gly355
Characteristics:	Recombinant Human IL-23R Protein is expressed from HEK293 with mFc (IgG1) tag at the C-Terminus.It contains Gly24-Gly355.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 μm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	Immobilized Human IL-23 alpha&Mouse IL-12, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Human IL-23R, mFc with the EC50 of 0.36µg/ml determined by ELISA. The affinity constant of 11.09 nM as determined in SPR assay (Biacore T200). See testing image for detail.

Target Details

Expiry Date:


12 months

. d. got 2 otdo	
Target:	IL23R
Alternative Name:	IL-23R (IL23R Products)
Background:	Interleukin 23 (IL-23) is a heterodimeric cytokine composed of two disulfide-linked subunits, a p19 subunit that is unique to IL-23, and a p40 subunit that is shared with IL-12. The functional IL-23 receptor complex consists of two receptor subunits, the IL-12 receptor beta 1 subunit (IL-12 R beta 1) and the IL-23-specific receptor subunit (IL-23 R). This receptor associates with IL12RB1 to form the interleukin-23 receptor. Binds IL23 and mediates T-cells, NK cells and possibly certain macrophage/myeloid cells stimulation probably through activation of the Jak-Stat signaling cascade.
Molecular Weight:	63.7 kDa. Due to glycosylation, the protein migrates to 80-115 kDa based on Tris-Bis PAGE result.
Pathways:	Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Activated T Cell Proliferation
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from $0.22\mu m$ filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Fi. Data	10




Surface Plasmon Resonance

Image 1. Human IL-23R, mFc Tag captured on CM5 Chip via Anti-Mouse Antibody can bind Human IL-23 alpha&IL-12 beta, His Tag with an affinity constant of 11.09 nM as determined in SPR assay (Biacore T200).



Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Human IL-23R is greater than 95 % as determined by SEC-HPLC.



SDS-PAGE

Image 3. Human IL-23R on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .

Please check the product details page for more images. Overall 4 images are available for ABIN7275031.