antibodies -online.com







LILRB1 Protein (AA 17-456) (Fc Tag)

Images



()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷


Quantity:	100 μg
Target:	LILRB1
Protein Characteristics:	AA 17-456
Origin:	Rhesus Monkey
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This LILRB1 protein is labelled with Fc Tag.

Product Details

Purpose:	Rhesus macaque LILRB1/CD85j/ILT2 Protein	
Sequence:	Ser17-His456	
Characteristics:	Recombinant Rhesus macaque LILRB1/CD85j/ILT2 Protein is expressed from HEK293 with hFc tag at the C-Terminus.It contains Ser17-His456.	
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC	
Sterility:	0.22 μm filtered	
Endotoxin Level:	Less than 1EU per μg by the LAL method.	
Biological Activity Comment:	gical Activity Comment: The affinity constant of 51.44 nM as determined in SPR assay (Biacore T200). See testing image for detail.	


Target Details

rarget betails		
Target:	LILRB1	
Alternative Name:	LILRB1 (LILRB1 Products)	
Background: Molecular Weight:	LILRB1, also known as CD85j and ILT2, is a 110 kDa transmembrane glycoprotein in the LILR immunoregulatory protein family. Mature human LILRB1 consists of a 438 amino acid (aa) extracellular domain (ECD) with 4 tandem Ig-like domains, a 21 aa transmembrane segment, and a 168 aa cytoplasmic domain with 4 inhibitory ITIM motifs.LILRB1 is a receptor for class I MHC antigens. Recognizes a broad spectrum of HLA-A, HLA-B, HLA-C, HLA-G and HLA-F alleles. Receptor for H301/UL18, a human cytomegalovirus class I MHC homolog. Ligand binding results in inhibitory signals and down-regulation of the immune response. 74.5 kDa. Due to glycosylation, the protein migrates to 80-90 kDa based on Tris-Bis PAGE results.	
UniProt:	F7H3G7	
Pathways:	Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response	
Application Details		
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water.	
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.	
Storage:	-20 °C,-80 °C	
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	
Expiry Date:	12 months	




Surface Plasmon Resonance

Image 1. Anti-LILRB1 Antibody immobilized on CM5 Chip can bind Rhesus macaque LILRB1, hFc Tag with an affinity constant of 51.44 nM as determined in SPR assay (Biacore T200).



Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Rhesus macaque LILRB1 is greater than 95 % as determined by SEC-HPLC.



SDS-PAGE

 $\label{limage 3.} \mbox{ Rhesus macaque LILRB1 on Tris-Bis PAGE under} \\ \mbox{reduced conditions. The purity is greater than 95 \% }.$