antibodies - online.com







MICA Protein (AA 24-308) (His-Avi Tag)

Images



Overview


Quantity:	100 μg
Target:	MICA
Protein Characteristics:	AA 24-308
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MICA protein is labelled with His-Avi Tag.

Product Details

Purpose:	Human MICA Protein
Sequence:	Glu24-Gln308
Characteristics:	Recombinant Human MICA Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.It contains Glu24-Gln308.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 μm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	Immobilized Human MICA, His Tag at 1µg/ml (100µl/Well) on the plate. Dose response curve for Human NKG2D, hFc Tag with the EC50 of 0.20µg/ml determined by ELISA. See testing image for detail.


Target Details

rarget Details	
Target:	MICA
Alternative Name:	MICA (MICA Products)
Background:	MICA (MHC class I chain-related gene A) is a transmembrane glycoprotein that functions as a ligand for human NKG2D. A closely related protein, MICB, shares 85 % amino acid identity with MICA. These proteins are distantly related to the MHC class I proteins. They possess three extracellular Ig-like domains, but they have no capacity to bind peptide or interact with beta 2-microglobulin
Molecular Weight:	35.8 kDa. Due to glycosylation, the protein migrates to 55-70 kDa based on Tris-Bis PAGE result.
UniProt:	Q96QC4
Pathways:	Activation of Innate immune Response, Transition Metal Ion Homeostasis
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months



Size-exclusion chromatography-High Pressure Liquid Chromatography


Image 1. The purity of Human MICA is greater than 95 % as determined by SEC-HPLC.



SDS-PAGE

Image 2. Human MICA on Tris-Bis PAGE under reduced condition. The purity is greater than $95\,\%$.

Human MICA, His Tag ELISA 0.1µg Human MICA, His Tag Per Well



ELISA

Image 3. Immobilized Human MICA, His Tag at 1 μ g/mL (100 μ L/Well) on the plate. Dose response curve for Human NKG2D, hFc Tag with the EC50 of 0.20 μ g/mL determined by ELISA.