antibodies

Datasheet for ABIN7275299 MUC16 Protein (AA 12660-12923) (His-Avi Tag)

4 Images



Overview


Quantity:	100 µg
Target:	MUC16 (CA125)
Protein Characteristics:	AA 12660-12923
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MUC16 protein is labelled with His-Avi Tag.

Product Details


Purpose:	Human CA125/MUC16 Protein
Sequence:	Gly12660-Met12923
Characteristics:	Recombinant Human CA125/MUC16 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.It contains Gly12660-Met12923.
Purity:	> 95 % as determined by Tris-Bis PAGE
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per μ g by the LAL method.
Biological Activity Comment:	Immobilized Human CA125, His Tag at 2µg/ml (100µl/Well) on the plate. Dose response curve for Human MSLN, hFc Tag with the EC50 of 2.8µg/ml determined by ELISA. See testing image
	for detail.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7275299 | 01/18/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Target:	MUC16 (CA125)
Alternative Name:	CA125 (CA125 Products)
Background:	MUC16, also known as the CA125 antigen, is a mucin protein that may be found in type I transmembrane or secreted forms that are used monitor the progress of epithelial ovarian cancer therapy.Thought to provide a protective, lubricating barrier against particles and infectious agents at mucosal surfaces.Binding to MSLN mediates heterotypic cell adhesion. This may contribute to the metastasis of ovarian cancer to the peritoneum by initiating cell attachment to the mesothelial epithelium via binding to MSLN.
Molecular Weight:	31.3 kDa. Due to glycosylation, the protein migrates to 50-80 kDa based on Tris-Bis PAGE result.
UniProt:	Q8WXI7
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months



Human CA125, His Tag ELISA 0.5µg Human CA125, His Tag Per Well



ELISA

ELISA

ELISA.

Image 3. Immobilized Human CA125 at $5 \mu g/mL$ (100 μ L/well) on the plate. Dose response curve for Human MSLN, hFc Tag with the EC50 of 1.14 $\mu g/mL$ determined by ELISA.

Please check the product details page for more images. Overall 4 images are available for ABIN7275299.

Image 1. Immobilized Human CA125, His Tag at $2 \mu g/mL$ (100 $\mu L/Well$) on the plate. Dose response curve for Human MSLN, hFc Tag with the EC50 of 2.8 $\mu g/mL$ determined by

Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Human CA125 is greater than 95 % as determined by SEC-HPLC.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7275299 | 01/18/2024 | Copyright antibodies-online. All rights reserved.