antibodies

Datasheet for ABIN7275632 SLAMF7 Protein (AA 23-226) (His-Avi Tag)





Overview

Quantity:	100 µg
Target:	SLAMF7
Protein Characteristics:	AA 23-226
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SLAMF7 protein is labelled with His-Avi Tag.


Product Details


Purpose:	Human SLAMF7/CRACC/CD319 Protein
Sequence:	Ser23-Met226
Characteristics:	Recombinant Human SLAMF7/CRACC/CD319 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.It contains Ser23-Met226.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per μ g by the LAL method.
Biological Activity Comment:	Immobilized Human SLAMF7, His Tag at 1μ g/ml (100 μ l/well) on the plate. Dose response curve
	for Anti-SLAMF7 Antibody, hFc Tag with the EC50 of 20.9ng/ml determined by ELISA. See
	testing image for detail.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7275632 | 01/18/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Target:	SLAMF7
Alternative Name:	SLAMF7 (SLAMF7 Products)
Background:	CD2-like receptor activating cytotoxic cells (CRACC), also known as CS1, novel Ly9, SLAMF7,
	and CD319, is a 65-75 kDa type I transmembrane glycoprotein in the SLAM subgroup of the
	CD2 family.Self-ligand receptor of the signaling lymphocytic activation molecule (SLAM) family.
	SLAM receptors triggered by homo- or heterotypic cell-cell interactions are modulating the
	activation and differentiation of a wide variety of immune cells and thus are involved in the
	regulation and interconnection of both innate and adaptive immune response.
Molecular Weight:	25.3 kDa. Due to glycosylation, the protein migrates to 40-55 kDa based on Tris-Bis PAGE result
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is
Reconstitution:	
Reconstitution: Buffer:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is
	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water.
	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water. Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as
Buffer:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water. Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Buffer: Storage:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water. Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization. -20 °C,-80 °C
Buffer: Storage:	 Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water. Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization. -20 °C,-80 °C -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after





Log Anti-SLAMF7 Antibody, hFc Tag Conc.(µg/ml)

SDS-PAGE

Image 1. Human SLAMF7 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .

Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Human SLAMF7 is greater than 95 % as determined by SEC-HPLC.

ELISA

Image 3. Immobilized Human SLAMF7, His Tag at $1 \mu g/mL$ (100 $\mu L/well$) on the plate. Dose response curve for Anti-SLAMF7 Antibody, hFc Tag with the EC50 of 20.9 ng/mL determined by ELISA.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7275632 | 01/18/2024 | Copyright antibodies-online. All rights reserved.