antibodies -online.com





Datasheet for ABIN7275726

TIGIT Protein (AA 22-141) (His tag)

3 Images



Go to Product page

Overview

Quantity:	100 μg
Target:	TIGIT
Protein Characteristics:	AA 22-141
Origin:	Cynomolgus, Rhesus Monkey
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TIGIT protein is labelled with His tag.

Product Details

Purpose:	Cynomolgus/Rhesus macaque TIGIT Protein
Sequence:	Met22-Ile141
Characteristics:	Recombinant Cynomolgus/Rhesus macaque TIGIT Protein is expressed from HEK293 with His tag at the C-Terminus.It contains Met22-Ile141.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per μg by the LAL method.
Biological Activity Comment:	Immobilized Cynomolgus TIGIT, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-TIGIT Antibody, hFc Tag with the EC50 of 30.3ng/ml determined by ELISA. See testing image for detail.


Target Details

Expiry Date:


12 months

Target Details	
Target:	TIGIT
Alternative Name:	TIGIT (TIGIT Products)
Background:	TIGIT, also called Vstm3, Vsig9, and WUCAM, is a transmembrane protein in the CD28 family of the Ig superfamily proteins.IGIT is expressed at low levels on peripheral memory and regulatory CD4 T-cells and NK cells and is up-regulated following activation of these cells.
Molecular Weight:	14.4 kDa. Due to glycosylation, the protein migrates to 18-25 kDa based on Tris-Bis PAGE result.
NCBI Accession:	XP_015300911
Pathways:	Cancer Immune Checkpoints
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Cynomolgus TIGIT, His Tag ELISA 0.05µg Cynomolgus TIGIT, His Tag Per Well



Log Anti-TIGIT Antibody, hFc Tag Conc.(µg/ml)



20 - 17.5 - 15 - 12.5 - 10 - 7.5 - 5 - 2.5 - 0

ELISA

Image 1. Immobilized Cynomolgus TIGIT, His Tag at $0.5\,\mu$ g/mL (100 μ L/Well) on the plate. Dose response curve for Anti-TIGIT Antibody, hFc Tag with the EC50 of 30.3 ng/mL determined by ELISA.

SDS-PAGE

Image 2. Cynomolgus TIGIT on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 3. The purity of Cynomolgus TIGIT is greater than 95 % as determined by SEC-HPLC.