antibodies

# Datasheet for ABIN7275778 TNFRSF10A Protein (His-Avi Tag,Biotin)





### Overview

Quantity:	100 µg
Target:	TNFRSF10A
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TNFRSF10A protein is labelled with His-Avi Tag,Biotin.

## Product Details

Sequence:	Pro34-Asn239
Purity:	> 95% as determined by Tris-Bis PAGE,> 95% as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per $\mu$ g by the LAL method.
Biological Activity Comment:	Immobilized Human TRAIL, No Tag at 5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Human TRAIL R1, His Tag with the EC50 of 0.31µg/ml determined by ELISA. The affinity constant of 0.315 nM as determined in SPR assay (Biacore T200). See testing image for detail.

## Target Details

Target:	TNFRSF10A
Alternative Name:	TRAIL R1 (TNFRSF10A Products)
Background:	DR4, CD261, TNFRSF10A, TRAIL-R, APO2, TRAIL R1, MGC9365, Tumor necrosis factor receptor

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7275778 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

## Target Details


	superfamily member 10A (TNFRSF10A) is also known as TNF-related apoptosis-inducing
	ligand receptor 1 (TRAIL-R1), Death receptor 4 (DR4), CD261 and APO2, which belongs to TNF
	superfamily.The expression of apoptosis-inducing TRAIL-R1 and TRAIL-R2 and of the decoy
	receptors TRAIL-R3 and TRAIL-R4 was systematically studied in all developmental stages of
	peripheral B cells isolated from the blood and secondary lymphoid organs. Expression of
	TRAIL-Rs is modulated along development, with highest levels observed in germinal center B
	cells.
Molecular Weight:	24.7 kDa. Due to glycosylation, the protein migrates to 25-30 kDa based on Tris-Bis PAGE result.
UniProt:	000220
Pathways:	Apoptosis, Positive Regulation of Endopeptidase Activity
Application Details	
Restrictions:	For Research Use only
Handling	
Handling Format:	Lyophilized
	Lyophilized Centrifuge tubes before opening. Reconstituting to a concentration more than 100 µg/mL is
Format:	
Format:	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 $\mu$ g/mL is
Format:	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 µg/mL is recommended (usually we use 1 mg/mL solution for lyophilization). Dissolve the lyophilized
Format: Reconstitution:	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 µg/mL is recommended (usually we use 1 mg/mL solution for lyophilization). Dissolve the lyophilized protein in distilled water.
Format: Reconstitution:	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 µg/mL is recommended (usually we use 1 mg/mL solution for lyophilization). Dissolve the lyophilized protein in distilled water. Lyophilized from 0.22µm filtered solution in PBS ( pH 7.4). Normally 5 % trehalose is added as
Format: Reconstitution: Buffer:	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 µg/mL is recommended (usually we use 1 mg/mL solution for lyophilization). Dissolve the lyophilized protein in distilled water. Lyophilized from 0.22µm filtered solution in PBS ( pH 7.4). Normally 5 % trehalose is added as protectant before lyophilization.
Format: Reconstitution: Buffer: Storage:	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 µg/mL is recommended (usually we use 1 mg/mL solution for lyophilization). Dissolve the lyophilized protein in distilled water. Lyophilized from 0.22µm filtered solution in PBS ( pH 7.4). Normally 5 % trehalose is added as protectant before lyophilization. 4 °C,-80 °C

mAU

30

20

15




## **ELISA**

Image 1. Immobilized Human TRAIL, No Tag at 5 µg/mL (100 µL/well) on the plate. Dose response curve for Biotinylated Human TRAIL R1, His Tag with the EC50 of 0.31 µg/mL determined by ELISA.

#### Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Biotinylated Human TRAIL R1 is greater than 95 % as determined by SEC-HPLC.



10KD

#### SDS-PAGE

Image 3. Biotinylated Human TRAIL R1 on Tris-Bis PAGE under reduced conditions. The purity is greater than 95 % .

Please check the product details page for more images. Overall 5 images are available for ABIN7275778.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7275778 | 09/10/2023 | Copyright antibodies-online. All rights reserved.