# antibodies - online.com







# TYRO3 Protein (AA 41-427) (His tag)

**Images** 



#### Overview

Quantity:	100 μg
Target:	TYRO3
Protein Characteristics:	AA 41-427
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TYRO3 protein is labelled with His tag.

#### **Product Details**


Purpose:	Cynomolgus TYRO3 Protein
Sequence:	Gly41-Ser427
Characteristics:	Recombinant Cynomolgus TYR03 Protein is expressed from HEK293 with His tag at the C-Terminus.It contains Gly41-Ser427.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by SEC-HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.

## **Target Details**

Target:	TYRO3
Alternative Name:	TYRO3 (TYRO3 Products)


## **Target Details**

O .	
Background:	The TAM receptors (Tyro3, Axl and MerTK) are promising therapeutic targets on tumorassociated macrophages. The TAM receptors are a family of receptor tyrosine kinases with shared ligands Gas6 and Protein S that skew macrophage polarization towards a pro-tumor M2-like phenotype. In macrophages, the TAM receptors also promote apoptotic cell clearance, a tumor-promoting process called efferocytosis. The TAM receptors bind the "eat-me" signal phosphatidylserine on apoptotic cell membranes using Gas6 and Protein S as bridging ligands.
Molecular Weight:	42.47 kDa. Due to glycosylation, the protein migrates to 50-70 kDa based on Tris-Bis PAGE result.
UniProt:	A0A2K5WCY8
Pathways:	RTK Signaling
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu$ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months



#### **SDS-PAGE**

**Image 1.** Cynomolgus TYRO3 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .



Size-exclusion chromatography-High Pressure Liquid Chromatography

**Image 2.** The purity of Cynomolgus TYRO3 is greater than 95 % as determined by SEC-HPLC.