



[Go to Product page](#)

Datasheet for ABIN7275832

## FLT4 Protein (AA 25-776) (Fc Tag)

### 3 Images

#### Overview

Quantity:	100 µg
Target:	FLT4
Protein Characteristics:	AA 25-776
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This FLT4 protein is labelled with Fc Tag.

#### Product Details

Purpose:	Human VEGF R3/FLT4 Protein
Sequence:	Tyr25-Ile776
Characteristics:	Recombinant Human VEGF R3/FLT4 Protein is expressed from HEK293 with hFc tag at the C-Terminus. It contains Tyr25-Ile776.
Purity:	> 95 % as determined by Tris-Bis PAGE, > 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	Immobilized Human VEGF-C, His Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Human VEGF R3, hFc Tag with the EC50 of 19.8ng/ml determined by ELISA. See testing image for detail.

## Target Details

---

Target:	FLT4
Alternative Name:	VEGF R3 ( <a href="#">FLT4 Products</a> )
Background:	Vascular endothelial growth factor (VEGF) and its receptors VEGF-R1, -R2 and -R3 play important roles in tumor angiogenesis and are associated with poor prognosis in several solid tumors. VEGF-R1, -R2 and -R3 were highly expressed in CRC cells and stromal vessels. VEGF-R1 strong positive staining correlated with shorter survival after CRC surgery.
Molecular Weight:	111.3 kDa. Due to glycosylation, the protein migrates to 135-155 kDa based on Tris-Bis PAGE result.
Pathways:	<a href="#">RTK Signaling</a> , <a href="#">Signaling Events mediated by VEGFR1 and VEGFR2</a> , <a href="#">VEGF Signaling</a>

## Application Details


---

Restrictions: For Research Use only

## Handling


---

Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in 50 mM Tris, 150 mM NaCl, 100 mM Glycine ( pH 7.5). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months



### SDS-PAGE


**Image 1.** Human VEGF R3/FLT4 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .



### Size-exclusion chromatography-High Pressure Liquid Chromatography

**Image 2.** The purity of Human VEGF R3/FLT4 is greater than 95 % as determined by SEC-HPLC.

**Human VEGF R3, hFc Tag ELISA**  
0.1µg Human VEGF-C, His Tag Per Well



### ELISA

**Image 3.** Immobilized Human VEGF-C, His Tag at 1 µg/mL (100 µL/well) on the plate. Dose response curve for Human VEGF R3, hFc Tag with the EC50 of 19.8 ng/mL determined by ELISA.