



[Go to Product page](#)

Datasheet for ABIN7278028  
**anti-Ly76 antibody (APC)**

1 Image

### Overview

Quantity:	100 µg
Target:	Ly76
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This Ly76 antibody is conjugated to APC
Application:	Flow Cytometry (FACS)

### Product Details

Clone:	TER-119
Isotype:	IgG2b kappa
Purification:	This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

### Target Details

Target:	Ly76
Alternative Name:	TER-119 ( <a href="#">Ly76 Products</a> )
Background:	The TER-119 antibody is named for the antigen to which it binds, a 52 kDa surface protein that is associated with glycophorin-A. TER-119 is considered to be a lineage marker for later stages

## Target Details

of erythroid cell development, as its expression begins at the pro-erythroblast stage. TER-119 antigen is not expressed at either BFU-E or CFU-E stages, i.e. prior to the pro-erythroblast stage.

Gene ID: 104231

## Application Details

**Application Notes:** This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indicated). The amount of antibody required for optimal staining of a cell sample should be determined empirically in your system.

**Comment:** 0.2 mg/mL

**Restrictions:** For Research Use only

## Handling

**Buffer:** 10 mM NaH<sub>2</sub>PO<sub>4</sub>, 150 mM NaCl, 0.09 % Sodium azide, 0.1 % gelatin, pH 7.2

**Preservative:** Sodium azide


**Precaution of Use:** This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

**Storage:** 4 °C

**Storage Comment:** 2-8°C protected from light

**Expiry Date:** 12 months

## Images



### Flow Cytometry

**Image 1.** C57Bl/6 bone marrow cells were stained with FITC Anti-Mouse CD45 (ABIN6961266) and 0.5 µg APC Anti-Mouse TER-119 (ABIN6961266) (right panel) or 0.5 µg APC Rat IgG2b isotype control (left panel).