

Datasheet for ABIN7278038

anti-Interferon gamma antibody (APC)**2** Images[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	Interferon gamma (IFNG)
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This Interferon gamma antibody is conjugated to APC
Application:	Flow Cytometry (FACS)

Product Details

Clone:	XMG1-2
Isotype:	IgG1 kappa
Purification:	This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

Target Details

Target:	Interferon gamma (IFNG)
Alternative Name:	IFN gamma (IFNG Products)
Background:	The XMG1.2 antibody is specific for mouse Interferon-gamma (IFN-g), a 20 kDa type II cytokine known for its central roles in protection against bacterial or viral pathogens and for its anti-

Target Details

tumor properties. IFN-g is secreted by several types of immune cells, which allow the cytokine to modulate innate immunity, when secreted by NK and NKT cells, and to function in support of adaptive immunity when secreted by Th1 and CD8+ T cells (CTLs). The XMG1.2 antibody is suitable for detection of intracellular IFN-g protein, e.g. by flow cytometry, as well as for quantitative analysis of the secreted protein by ELISA, when paired with an appropriate secondary antibody. This clone is also widely used for neutralization of the functional activity of IFN-g in a variety of assays.

Gene ID: 15978

UniProt: [P01580](#)

Pathways: [Interferon-gamma Pathway](#), [Cellular Response to Molecule of Bacterial Origin](#), [Regulation of Leukocyte Mediated Immunity](#), [Positive Regulation of Immune Effector Process](#), [Production of Molecular Mediator of Immune Response](#), [ER-Nucleus Signaling](#), [Regulation of Carbohydrate Metabolic Process](#), [Protein targeting to Nucleus](#), [Autophagy](#)

Application Details

Application Notes: This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indicated). The amount of antibody required for optimal staining of a cell sample should be determined empirically in your system.

Comment: 0.2 mg/mL

Restrictions: For Research Use only

Handling

Buffer: 10 mM NaH₂PO₄, 150 mM NaCl, 0.09 % Sodium azide, 0.1 % gelatin, pH 7.2


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C


Storage Comment: 2-8°C protected from light

Expiry Date: 12 months



Flow Cytometry

Image 1. C57Bl/6 splenocytes were stimulated and stained with FITC Anti-Mouse CD3e (ABIN6961277), followed by intracellular staining with 0.06 μ g APC Anti-Mouse IFN gamma (XMG1.2) manufactured by antibodies-online (left panel) or eBioscience (right panel).



Flow Cytometry

Image 2. C57Bl/6 splenocytes were stimulated with PMA and Ionomycin (right panel) or unstimulated (left panel) and then stained with FITC Anti-Mouse CD3e (ABIN6961277), followed by intracellular staining with 0.06 μ g APC Anti-Mouse IFN gamma (ABIN6961277).