

#### Datasheet for ABIN7278328

## anti-CSF1R antibody (PE)

# 1 Image



Go to Product page

()	ve	rvi	0	W
$\circ$	v C	1 V I	$\sim$	v v

Quantity:	100 μg
Target:	CSF1R
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This CSF1R antibody is conjugated to PE
Application:	Flow Cytometry (FACS)

#### **Product Details**

AFS98

Clone:

Isotype:	IgG2a kappa
Purification:	This monoclonal antibody was purified from tissue culture supernatant via affinity
	chromatography. The purified antibody was conjugated under optimal conditions, with
	unreacted dye removed from the preparation. It is recommended to store the product undiluted
	at 4°C, and protected from prolonged exposure to light. Do not freeze.

#### **Target Details**

Target:	CSF1R
Alternative Name:	CD115 (c-fms) (CSF1R Products)
Background:	The AFS98 antibody is specific for mouse CD115, also known as Colony-Stimulating Factor-1 Receptor (CSF-1R), a 145 kDa receptor from the PDGF receptor family. Receptor activation by

the ligands IL-34 or CSF-1 (M-CSF) occurs via homodimerization of CD115 and subsequent
tyrosine phosphorylation and ubiquitination of intracellular domains. CD115 signaling promotes
differentiation of myeloid pre- cursors, as well as the continued regulation of proliferation,
survival and function of mononuclear phagocytes, dendritic cells and osteo- clasts. While IL-34
and CSF-1 may induce similar cellular responses, they are differentially expressed and as such
exert complimentary actions via CD115.The AFS98 antibody may be used for identification of
myeloid lineage cells by flow cytometry, and is commonly used for in vivo or in vitro
neutralization of CSF-1 Receptor.
12978
P09581

UniProt:	P0958

Pathways: RTK Signaling, Inositol Metabolic Process, Cell-Cell Junction Organization

### **Application Details**

Gene ID:


Application Notes:	This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells,
	or an appropriate cell type (where indi-cated). The amount of antibody required for optimal
	staining of a cell sample should be determined empirically in your system.

Comment:	0.2 mg/mL

Restrictions: For Research Use only

#### Handling

Buffer:	10 mM NaH2PO4, 150 mM NaCl, 0.09 % Sodium azide, 0.1 % gelatin, pH 7.2
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	2-8°C protected from light
Expiry Date:	12 months



#### Flow Cytometry

**Image 1.** C57BI/6 peripheral blood cells were stained with APC Anti-Mouse CD11b (ABIN6961534) and 0.125  $\mu$ g PE Anti-Mouse CD115 (ABIN6961534) (right panel) or 0.125  $\mu$ g PE Rat IgG2a (left panel).