

Datasheet for ABIN7278360

anti-Interferon gamma antibody (PE)

XMG1-2



Overview

Quantity:	100 μg
Target:	Interferon gamma (IFNG)
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This Interferon gamma antibody is conjugated to PE
Application:	Flow Cytometry (FACS)

Product Details

Clone:

Isotype:	IgG1 kappa
Purification:	This monoclonal antibody was purified from tissue culture supernatant via affinity
	chromatography. The purified antibody was conjugated under optimal conditions, with
	unreacted dye removed from the preparation. It is recommended to store the product undiluted
	at 4°C, and protected from prolonged exposure to light. Do not freeze.

Target Details

Target:	Interferon gamma (IFNG)
Alternative Name:	IFN gamma (IFNG Products)
Background:	The XMG1.2 antibody is specific for mouse Interferon-gamma (IFN-g), a 20 kDa type II cytokine
	known for its central roles in protection against bacterial or viral pathogens and for its anti-

IFN-g in a variety of assays.
secondary antibody. This clone is also widely used for neutralization of the functional activity of
quantitative analysis of the secreted protein by ELISA, when paired with an appropriate
suitable for detection of intracellular IFN-g protein, e.g. by flow cytometry, as well as for
adaptive immunity when secreted by Th1 and CD8+ T cells (CTLs). The XMG1.2 antibody is
to modulate innate immunity, when secreted by NK and NKT cells, and to function in support of
tumor properties. IFN-g is secreted by several types of immune cells, which allow the cytokine

O ID:	1 5070
Gene ID:	15978

UniProt: P01580

Interferon-gamma Pathway, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, ER-Nucleus Signaling, Regulation of Carbohydrate Metabolic Process, Protein targeting to Nucleus, Autophagy

Application Details

Pathways:

Application Notes:	This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells,
	or an appropriate cell type (where indi-cated). The amount of antibody required for optimal
	staining of a cell sample should be determined empirically in your system

Comment: 0.2 mg/mL

Restrictions: For Research Use only

Handling

Buffer:	10 mM NaH2PO4, 150 mM NaCl, 0.09 % Sodium azide, 0.1 % gelatin, pH 7.2
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	2-8°C protected from light
Expiry Date:	12 months