

Datasheet for ABIN7278425
anti-IL7R antibody (PE-Cy7)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	IL7R
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This IL7R antibody is conjugated to PE-Cy7
Application:	Flow Cytometry (FACS)

Product Details

Clone:	A7R34
Isotype:	IgG2a kappa
Purification:	This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

Target Details

Target:	IL7R
Alternative Name:	CD127 (IL-7Ra) (IL7R Products)
Background:	The A7R34 antibody is specific for mouse CD127, a 60-90 kDa cell surface protein also known as the Interleukin-7 Receptor alpha chain, or IL-7R alpha. CD127 is typically expressed at the

Target Details

cell surface as a heterodimer with the common gamma chain (CD132). This complex acts as the functional receptor for IL-7, a cytokine important in T and B cell development, and in mature T cell homeostasis. A second cytokine known as Thymic Stromal Lymphopoietin (TSLP) also binds to a receptor complex of CD127 and the TSLPR chain to trigger activation of dendritic cells, and is involved in B cell development, allergy and autoimmunity. The A7R34 antibody may be used as a phenotypic marker for CD127 on immature B cells, on subsets of thymocytes which are double negative (CD4-CD8-) or single positive (CD4+ or CD8+), and at low levels on mature, peripheral T cells. CD127 is a key marker, when used in combination with CD4 and CD25, to distinguish Treg and effector/memory Treg populations known as T(REM).


Gene ID:	16172
UniProt:	P16872
Pathways:	JAK-STAT Signaling , Regulation of Leukocyte Mediated Immunity , Production of Molecular Mediator of Immune Response , Regulation of Cell Size

Application Details

Application Notes:	This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indicated). The amount of antibody required for optimal staining of a cell sample should be determined empirically in your system.
Comment:	0.2 mg/mL
Restrictions:	For Research Use only

Handling

Buffer:	10 mM NaH ₂ PO ₄ , 150 mM NaCl, 0.09 % Sodium azide, 0.1 % gelatin, pH 7.2
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	2-8°C protected from light
Expiry Date:	12 months



Flow Cytometry

Image 1. C57Bl/6 splenocytes were stained with FITC Anti-Mouse CD3 (ABIN6961635) and 0.5 µg PE-Cy7 Anti-Mouse CD127 (ABIN6961635) (right panel) or 0.5 µg PE-Cy7 Rat IgG2a isotype control (left panel).