

# Datasheet for ABIN7278434

# anti-Interferon gamma antibody (PE-Cy7)

XMG1-2



#### Overview

Quantity:	100 μg
Target:	Interferon gamma (IFNG)
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This Interferon gamma antibody is conjugated to PE-Cy7
Application:	Flow Cytometry (FACS)

#### **Product Details**

Clone:

Isotype:	IgG1 kappa
Purification:	This monoclonal antibody was purified from tissue culture supernatant via affinity
	chromatography. The purified antibody was conjugated under optimal conditions, with
	unreacted dye removed from the preparation. It is recommended to store the product undiluted
	at 4°C, and protected from prolonged exposure to light. Do not freeze.

# **Target Details**

Target:	Interferon gamma (IFNG)
Alternative Name:	IFN gamma (IFNG Products)
Background:	The XMG1.2 antibody is specific for mouse Interferon-gamma (IFN-g), a 20 kDa type II cytokine
	known for its central roles in protection against bacterial or viral pathogens and for its anti-

tumor properties. IFN-g is secreted by several types of immune cells, which allow the cytokine
to modulate innate immunity, when secreted by NK and NKT cells, and to function in support of
adaptive immunity when secreted by Th1 and CD8+ T cells (CTLs). The XMG1.2 antibody is
suitable for detection of intracellular IFN-g protein, e.g. by flow cytometry, as well as for
quantitative analysis of the secreted protein by ELISA, when paired with an appropriate
secondary antibody. This clone is also widely used for neutralization of the functional activity of
IFN-g in a variety of assays.

Gene ID: 15978

UniProt: P01580

Pathways: Interferon-gamma Pathway, Cellular Response to Molecule of Bacterial Origin, Regulation of

Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, ER-Nucleus Signaling, Regulation of Carbohydrate

Metabolic Process, Protein targeting to Nucleus, Autophagy

### **Application Details**

Application Notes: This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indi- cated). The amount of antibody required for optimal

staining of a cell sample should be determined empirically in your system.

Comment: 0.2 mg/mL

Restrictions: For Research Use only

# Handling

Buffer:	10 mM NaH2PO4, 150 mM NaCl, 0.09 % Sodium azide, 0.1 % gelatin, pH 7.2
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C

Storage Comment: 2-8°C protected from light

Expiry Date: 12 months