

Datasheet for ABIN7278474

anti-L-Selectin antibody (PerCP-Cy5.5)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	L-Selectin (SELL)
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This L-Selectin antibody is conjugated to PerCP-Cy5.5
Application:	Flow Cytometry (FACS)

Product Details

Clone:	MEL-14
Isotype:	IgG2a kappa
Purification:	This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

Target Details

Target:	L-Selectin (SELL)
Alternative Name:	CD62L (L-Selectin) (SELL Products)
Background:	The MEL-14 antibody is specific for mouse CD62L, also known as L-Selectin, a cell adhesion molecule which facilitates lymphocyte "rolling" on activated vascular endothelium and homing

Target Details

to high endothelial venules (HEV) as immune cells transmigrate from blood into peripheral tissues. L-Selectin is a member of a family of Selectin molecules which act together with the integrin family of adhesion molecules to mediate leukocyte-endothelial interactions. L-Selectin is characteristically expressed by neutrophils, and is also found on B cells, monocytes, granulocytes, and at varying levels on naive, effector and memory T cells. It is rapidly shed upon cell activation, releasing into the circulation a soluble form whose biological role is of particular interest in cancer biology research. The MEL -14 antibody may be used as a phenotypic marker for CD62L expression on a variety of immune cell types. Please note that CD62L (L-Selectin) itself is also referred to as MEL-14 in the literature.

Gene ID: 20343

UniProt: [P18337](#)

Application Details

Application Notes: This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indicated). Please refer to the figure legend for the optimal concentration used to stain the tissue shown. We recommend titrating the antibody under your specific conditions to determine the optimal concentration of antibody needed in your experimental system.

Comment: 0.2 mg/mL

Restrictions: For Research Use only

Handling

Buffer: 10 mM NaH₂PO₄, 150 mM NaCl, 0.09 % Sodium azide, 0.1 % gelatin, pH 7.2

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C

Storage Comment: 2-8°C protected from light

Expiry Date: 12 months