

Datasheet for ABIN7278559

anti-CD45 antibody (violetFluor™ 450)

2 Images



Go to Product page

Overview

Quantity:	100 μg
Target:	CD45 (PTPRC)
Reactivity:	Human, Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This CD45 antibody is conjugated to violetFluor™ 450
Application:	Flow Cytometry (FACS)

Product Details

Clone:	RA3-6B2
Isotype:	IgG2a kappa
Cross-Reactivity:	Cat
Purification:	This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

Target Details

Target:	CD45 (PTPRC)
Alternative Name:	CD45R (B220) (PTPRC Products)
Background:	The RA3-6B2 antibody reacts with the human and mouse CD45 isoform known as CD45R, or

B220, a protein tyrosine phosphatase of 220 kDa. CD45 is one of the most abundant hematopoietic markers, and is expressed on all leukocytes (the Leukocyte Common Antigen, LCA). Various isoforms are generated and expressed in cell-specific patterns, all critical for leukocyte function. In mouse, the CD45R/B220 isoform is predominantly found on B cells, at varying levels on all stages from pro-B cells to activated B cells, and may also be detected on certain T cell and NK cell subsets. It is of note that B220 is not similarly expressed on human B cells, where it appears to be differentiation-specific and therefore expressed on only some B cell subsets. Other forms of CD45 with restricted cellular expression include CD45RA, CD45RB, CD45RO and several others. The RA3-6B2 antibody is one of the most consistently used leukocyte markers for B cells, T cell subsets and NK cell subsets in human and mouse. This antibody has also been reported as cross-reactive with feline CD45R/B220.

Gene ID: 19264, 5788

UniProt: P08575, P06800

Pathways: TCR Signaling, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune

Effector Process, Production of Molecular Mediator of Immune Response, CXCR4-mediated

Signaling Events, BCR Signaling

Application Details

Application Notes:

This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indi- cated). The amount of antibody required for optimal staining of a cell sample should be determined empirically in your system. violetFluor™ 450 dye is excited by the violet (405 nm) laser and has a peak emission of 450 nm. The most common band pass filters for this dye are 440/40 or 450/50. violetFluor™ 450 can be used as an alternative for Pacific Blue®, BD Horizon™ V450 or eFluor® 450.

Comment: 0.2 mg/mL

Restrictions: For Research Use only


Handling


Buffer:	10 mM NaH2PO4, 150 mM NaCl, 0.09 % Sodium azide, 0.1 % gelatin, pH 7.2
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.

Handling

Storage:	4 °C
Storage Comment:	2-8°C protected from light
Expiry Date:	12 months


Images





Flow Cytometry

Image 1. C57BI/6 splenocytes were stained with violetFluor 450 Anti-Mouse CD45R (B220) (RA3-6B2) manufactured by antibodies-online (left panel) or Horizon V450 Anti-Mouse CD45R (B220) (RA3-6B2) manufactured by BD Biosciences (right panel).



Hu/Mo CD45R (B220) (RA3-6B2) violetFluor 450

Flow Cytometry

Image 2. C57BI/6 splenocytes were stained with $0.5 \,\mu g$ violetFluor 450 Anti-Hu/Mo CD45R (B220) (ABIN6961807) (solid line) or $0.5 \,\mu g$ violetFluor 450 Rat IgG2a isotype control (dashed line).