antibodies - online.com









Images



Go to	Prod	uct	nage

$\overline{}$			
()	V/P	r\/	i٩٧٨

Quantity:	100 μL
Target:	Fascin (FSCN1)
Binding Specificity:	AA 11-110
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Fascin antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details


Immunogen:	KLH conjugated synthetic peptide derived from human Fascin1
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Pig
Purification:	Purified by Protein A.

Target Details

Torget	Faccin (FCCN1)	
Target:	Fascin (FSCN1)	


Target Details

raiget Details	
Alternative Name:	Fascin1 (FSCN1 Products)
Background:	Synonyms: HSN, SNL, p55, FAN1, Fascin, 55 kDa actin-bundling protein, Singed-like protein,
	FSCN1
	Background: Organizes filamentous actin into bundles with a minimum of 4.1:1 actin/fascin
	ratio. Plays a role in the organization of actin filament bundles and the formation of
	microspikes, membrane ruffles, and stress fibers. Important for the formation of a diverse set
	of cell protrusions, such as filopodia, and for cell motility and migration.
Gene ID:	6624
UniProt:	Q16658
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months




Immunofluorescence

Image 1. Formalin-fixed and paraffin embedded human gastric carcinoma with Anti-Fascin1 Polyclonal Antibody, Unconjugated (ABIN728293) at 1:300, followed by conjugation to the secondary antibody Goat Anti-Rabbit IgG, PE conjugated used at 1:200 dilution for 40 minutes at 37°C.



Immunohistochemistry

Image 2. Formalin-fixed and paraffin embedded rat brainstem with Anti-Fascin1 Polyclonal Antibody, Unconjugated (ABIN728293) at 1:300, followed by conjugation to the secondary antibody and DAB staining



Immunofluorescence (Paraffin-embedded Sections)

Image 3. Formalin-fixed and paraffin embedded human gastric carcinoma with Anti-Fascin1 Polyclonal Antibody, Unconjugated at 1:300, followed by conjugation to the secondary antibody Goat Anti-Rabbit IgG, PE conjugated - PE)used at 1:200 dilution for 40 minutes at 37°C.

Please check the product details page for more images. Overall 5 images are available for ABIN728293.