antibodies - online.com







anti-Collagen IV antibody (AA 1401-1500)





Overview

Quantity:	100 μL
Target:	Collagen IV (COL4)
Binding Specificity:	AA 1401-1500
Reactivity:	Human, Mouse, Rat, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Collagen IV antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from mouse Collagen alpha-1(IV) chain
Isotype:	IgG
Specificity:	This antibody has 79 % cross reactivity with Collagen alpha-5(IV) chain.
Cross-Reactivity:	Human, Mouse, Rabbit, Rat
Predicted Reactivity:	Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	Collagen IV (COL4)
Alternative Name:	Collagen 4 (COL4 Products)
Background:	Synonyms: Bru, Raw, Svc, Col4a-1, Del(8)44H, Collagen alpha-1(IV) chain, Col4a1
	Background: Type IV collagen is the major structural component of glomerular basement
	membranes (GBM), forming a 'chicken-wire' meshwork together with laminins, proteoglycans
	and entactin/nidogen. Arresten, comprising the C-terminal NC1 domain, inhibits angiogenesis
	and tumor formation. The C-terminal half is found to possess the anti-angiogenic activity.
	Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits
	expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for
	alpha1/beta1 integrin (By similarity).
Gene ID:	12826
UniProt:	P02463
Application Details	

Restrictions:	For Research Use only
	IF(ICC) 1:50-200
	IF(IHC-F) 1:50-200
	IF(IHC-P) 1:50-200
	IHC-F 1:100-500
	IHC-P 1:200-400
	FCM 1:20-100
	ELISA 1:500-1000
Application Notes:	WB 1:300-5000


Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling


Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Images




Immunohistochemistry

Image 1. Formalin-fixed and paraffin embedded rat liver tissue with Anti-Collagen IV Polyclonal Antibody, Unconjugated (ABIN728413) at 1:200, followed by conjugation to the secondary antibody and DAB staining



Immunohistochemistry

Image 2. Formalin-fixed and paraffin embedded rabbit cartilage tissue with Anti-Collagen IV Polyclonal Antibody, Unconjugated (ABIN728413) at 1:200, followed by conjugation to the secondary antibody and DAB staining



Immunohistochemistry (Paraffin-embedded Sections)

Image 3. Formalin-fixed and paraffin embedded rabbit cartilage tissue with Anti-Collagen IV Polyclonal Antibody, Unconjugated at 1:200, followed by conjugation to the secondary antibody and DAB staining

Please check the product details page for more images. Overall 4 images are available for ABIN728413.	