## antibodies .- online.com





## anti-TTF1 antibody (AA 201-300) (FITC)



$\sim$		D	4	page
1-0	$T \cap$	$Pr \cap C$	шет	nane
$\cup$	LO.	1 100	IUCL	Dauc

/ N	11/0	K /	$\sim$	A /
	ve	1 \/	ı⊢۱	$\Lambda$
$\sim$	' V C	1 V		v v

Quantity:	100 μL
Target:	TTF1
Binding Specificity:	AA 201-300
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TTF1 antibody is conjugated to FITC
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human TTF-1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

## Target Details

Target:	TTF1
Alternative Name:	TTF1 (TTF1 Products)
Background:	Synonyms: BCH, BHC, NK-2, TEBP, TTF1, NKX2A, T/EBP, TITF1, TTF-1, NKX2.1, Homeobox
	protein Nkx-2.1, Homeobox protein NK-2 homolog A, Thyroid nuclear factor 1, Thyroid

	transcription factor 1, Thyroid-specific enhancer-binding protein, NKX2-1
	Background: Transcription factor that binds and activates the promoter of thyroid specific
	genes such as thyroglobulin, thyroperoxidase, and thyrotropin receptor. Crucial in the
	maintenance of the thyroid differentiation phenotype. May play a role in lung development and
	surfactant homeostasis. Activates the transcription of GNRHR and plays a role in enhancing the
	circadian oscillation of its gene expression. Represses the transcription of the circadian
	transcriptional repressor NR1D1 (By similarity).
Gene ID:	7080
UniProt:	P43699
Pathways:	Thyroid Hormone Synthesis, Regulation of Systemic Arterial Blood Pressure by Hormones,
	Feeding Behaviour
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months