antibodies - online.com







anti-RHOH antibody (C-Term)



Characteristics:

Purification:



Go	+0	Dr	\sim d	110	t n	00	
(7()	1()	-	()(1100		\mathbf{a}	ı

Overview	
Quantity:	100 μL
Target:	RHOH
Binding Specificity:	C-Term
Reactivity:	Human, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RHOH antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human RhoH.
Specificity:	Recognizes endogenous levels of RhoH protein.

Target Details	
Target:	RHOH
Alternative Name:	RhoH (RHOH Products)

The antibody was purified by immunogen affinity chromatography.

Rabbit polyclonal antibody to RhoH

Background: ARHH, TTF, Rho-related GTP-binding protein RhoH, GTP-binding protein TTF, Translocation

Target Details

	three four protein	
Gene ID:	399	
UniProt:	Q15669	


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)	
Restrictions:	For Research Use only	

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of RhoH staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of RhoH expression in HEK293T (A), HepG2 (B), Jurkat (C) whole cell lysates.