

Datasheet for ABIN7295004
anti-Bcl-2 antibody (pSer69)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	Bcl-2 (BCL2)
Binding Specificity:	pSer69
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Bcl-2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human BCL2.
Specificity:	Recognizes endogenous levels of BCL2 (pT69) protein.
Characteristics:	Rabbit polyclonal antibody to BCL2 (pT69)
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	Bcl-2 (BCL2)
Alternative Name:	BCL2 (BCL2 Products)
Background:	Apoptosis regulator Bcl-2

Target Details


Gene ID:	596, 12043, 24224
UniProt:	P10415 , P10417 , P49950
Pathways:	MAPK Signaling , PI3K-Akt Signaling , Apoptosis , Caspase Cascade in Apoptosis , Regulation of Muscle Cell Differentiation , Cell-Cell Junction Organization , Skeletal Muscle Fiber Development , Autophagy , Smooth Muscle Cell Migration , Negative Regulation of intrinsic apoptotic Signaling

Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of BCL2 (pT69) expression in HepG2 UV-treated (A), HeLa (B), HL60 (C), THP1 (D), Jurkat paclitaxel-treated (E), NIH3T3 H2O2-treated (F), PC12 H2O2-treated (G) whole cell lysates.



Immunohistochemistry

Image 2. Immunohistochemical analysis of BCL2 (pT69) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.