

# Datasheet for ABIN7295126

# anti-Caspase 9 antibody (Center)

3 Images



Go to Product page

_				
()	ve.	rv/	101	Λ

0.101.1011		
Quantity:	100 μL	
Target:	Caspase 9 (CASP9)	
Binding Specificity:	Center	
Reactivity:	Human, Monkey	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Caspase 9 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF),	
	Immunochromatography (IC)	
Product Details		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human	
	Caspase 9.	
Specificity:	Recognizes endogenous levels of Caspase 9 protein.	
Characteristics:	Rabbit polyclonal antibody to Caspase 9	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Target Details		
Target:	Caspase 9 (CASP9)	
Alternative Name:	Caspase 9 (CASP9 Products)	

### **Target Details**


Background:	MCH6, Caspase-9, CASP-9, Apoptotic protease Mch-6, Apoptotic protease-activating factor 3, APAF-3, ICE-like apoptotic protease 6, ICE-LAP6
Gene ID:	842
UniProt:	P55211
Pathways:	MAPK Signaling, RTK Signaling, Apoptosis, Caspase Cascade in Apoptosis, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Positive Regulation of Endopeptidase Activity


### **Application Details**


Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months







#### **Immunofluorescence**

Image 1. Immunofluorescent analysis of Caspase 9 staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

### Immunohistochemistry

**Image 2.** Immunohistochemical analysis of Caspase 9 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **Western Blotting**

**Image 3.** Western blot analysis of Caspase 9 expression in HeLa (A), NIH3T3 (B) whole cell lysates.