

Datasheet for ABIN7295160
anti-Cyclin F antibody (C-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	Cyclin F (CCNF)
Binding Specificity:	C-Term
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Cyclin F antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cyclin F.
Specificity:	Recognizes endogenous levels of Cyclin F protein.
Characteristics:	Rabbit polyclonal antibody to Cyclin F
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	Cyclin F (CCNF)
Alternative Name:	Cyclin F (CCNF Products)
Background:	FBX1, FBX01, Cyclin-F, F-box only protein 1

Target Details

Gene ID:	899, 12449, 117524
UniProt:	P41002 , P51944 , Q8K4F8


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Western Blotting

Image 1. Western blot analysis of Cyclin F expression in HEK293T (A), mouse liver (B), rat liver (C) whole cell lysates.



Immunohistochemistry

Image 2. Immunohistochemical analysis of Cyclin F staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.