# antibodies -online.com







# anti-CDC27 antibody (Center)





$C \circ t \circ$	Product	nogo
GU LU	FIGUUCI	Luaue

Overview	
Quantity:	100 μL
Target:	CDC27
Binding Specificity:	Center
Reactivity:	Human, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDC27 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CDC27.
Specificity:	Recognizes endogenous levels of CDC27 protein.
Characteristics:	Rabbit polyclonal antibody to CDC27
Purification:	The antibody was purified by immunogen affinity chromatography.
Target Details	
Target:	CDC27
Alternative Name:	CDC27 (CDC27 Products)
Background:	ANAPC3, D0S1430E, D17S978E, Cell division cycle protein 27 homolog, Anaphase-promoting

### **Target Details**

	complex subunit 3, APC3, CDC27 homolog, CDC27Hs, H-NUC
Gene ID:	996
UniProt:	P30260


### **Application Details**

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)
Restrictions:	For Research Use only

### Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

#### **Images**



## **Immunohistochemistry**

Image 1. Immunohistochemical analysis of CDC27 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



#### **Western Blotting**

Image 2. Western blot analysis of CDC27 expression in A549 (A), K562 (B), Jurkat (C), HEK293T (D), HeLa (E) whole cell lysates.