

Datasheet for ABIN7295334
anti-COMT antibody (Center)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	COMT
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This COMT antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human COMT.
Specificity:	Recognizes endogenous levels of COMT protein.
Characteristics:	Rabbit polyclonal antibody to COMT
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	COMT
Alternative Name:	COMT (COMT Products)
Background:	Catechol O-methyltransferase

Target Details

Gene ID:	1312, 12846, 24267
UniProt:	P21964 , O88587 , P22734
Pathways:	Steroid Hormone Biosynthesis , SARS-CoV-2 Protein Interactome


Application Details

Application Notes:	WB (1:500 - 1:1000), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence

Image 1. Immunofluorescent analysis of COMT staining in Raw264.7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell



nuclei (blue).

Western Blotting

Image 2. Western blot analysis of COMT expression in HEK293T (A), Raw264.7 (B), rat liver (C) whole cell lysates.