



[Go to Product page](#)

Datasheet for ABIN7295392  
**anti-CSR1 antibody (Center)**

2 Images

Overview

Quantity:	100 µL
Target:	CSR1
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat, Chicken
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CSR1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CSR1.
Specificity:	Recognizes endogenous levels of CSR1 protein.
Characteristics:	Rabbit polyclonal antibody to CSR1
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	CSR1
Alternative Name:	CSR1 ( <a href="#">CSR1 Products</a> )
Background:	CSR1, CYRP, Cysteine and glycine-rich protein 1, Cysteine-rich protein 1, CRP, CRP1, Epididymis

## Target Details

luminal protein 141, HEL-141

Gene ID: 1465, 13007, 29276

UniProt: [P21291](#), [P97315](#), [P47875](#)

## Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

## Images



### Western Blotting

**Image 1.** Western blot analysis of CSRP1 expression in Jurkat (A), HeLa (B), PC3 (C), DU145 (D), A10 (E) whole cell lysates.



### Immunohistochemistry

**Image 2.** Immunohistochemical analysis of CSRP1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.