

Datasheet for ABIN7295890
anti-HNRNPH1 antibody (Center)

2 Images

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	HNRNPH1
Binding Specificity:	Center
Reactivity:	Human, Cow, Chicken
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HNRNPH1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human hnRNP H.
Specificity:	Recognizes endogenous levels of hnRNP H protein.
Characteristics:	Rabbit polyclonal antibody to hnRNP H
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	HNRNPH1
Alternative Name:	HnRNP H (HNRNPH1 Products)
Background:	HNRNPH1, HNRPH, HNRPH1, Heterogeneous nuclear ribonucleoprotein H, hnRNP H,

Target Details

	HNRNPH2, FTP3, HNRPH2, Heterogeneous nuclear ribonucleoprotein H2, hnRNP H2, FTP-3, Heterogeneous nuclear ribonucleoprotein H', hnRNP H'
Gene ID:	3188
UniProt:	P31943 , P55795


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of hnRNP H staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with

haematoxylin and mounted with DPX.

Western Blotting

Image 2. Western blot analysis of hnRNP H expression in A549 (A), HeLa (B) whole cell lysates.

