

# Datasheet for ABIN7296136 anti-Keratin 10 antibody (Center)





Go to Product page

$\sim$			
( )\	<b>/</b> e	rVI	iew

Quantity:	100 μL	
Target:	Keratin 10 (KRT10)	
Binding Specificity:	Center	
Reactivity:	Human, Mouse, Rat, Dog, Rabbit	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Keratin 10 antibody is un-conjugated	
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunofluorescence (IF),	
	Immunochromatography (IC)	
Product Details		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human	
	Cytokeratin 10.	
Specificity:	Recognizes endogenous levels of Cytokeratin 10 protein.	
Characteristics:	Rabbit polyclonal antibody to Cytokeratin 10	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Target Details		
Target:	Keratin 10 (KRT10)	
Alternative Name:	Cytokeratin 10 (KRT10 Products)	

#### **Target Details**

Background:	KPP, Keratin, type I cytoskeletal 10, Cytokeratin-10, CK-10, Keratin-10, K10
Gene ID:	3858, 16661, 450225
UniProt:	P13645, P02535, Q6IFW6

## **Application Details**

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.	
Preservative:	Sodium azide	

should be handled by trained staff only.


Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

Expiry Date: 12 months

## **Images**


Precaution of Use:




# **Western Blotting**

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

**Image 1.** Western blot analysis of Cytokeratin 10 expression in HeLa (A), A431 (B), Raw264.7 (C), NIH3T3 (D), rat brain (E) whole cell lysates.





#### **Immunofluorescence**

Image 2. Immunofluorescent analysis of Cytokeratin 10 staining in A431 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

#### **Immunohistochemistry**

Image 3. Immunohistochemical analysis of Cytokeratin 10 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.