

Datasheet for ABIN7296144  
**anti-KRT17 antibody (C-Term)**



[Go to Product page](#)

3 Images

## Overview

Quantity:	100 µL
Target:	KRT17
Binding Specificity:	C-Term
Reactivity:	Human, Rat, Dog, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KRT17 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

## Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cytokeratin 17.
Specificity:	Recognizes endogenous levels of Cytokeratin 17 protein.
Characteristics:	Rabbit polyclonal antibody to Cytokeratin 17
Purification:	The antibody was purified by immunogen affinity chromatography.

## Target Details

Target:	KRT17
Alternative Name:	Cytokeratin 17 ( <a href="#">KRT17 Products</a> )

## Target Details

Background:	Keratin, type I cytoskeletal 17, 39.1, Cytokeratin-17, CK-17, Keratin-17, K17
Gene ID:	3872, 287702
UniProt:	<a href="#">Q04695</a> , <a href="#">Q6IFU8</a>


## Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

## Images



### Immunofluorescence


**Image 1.** Immunofluorescent analysis of Cytokeratin 17 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI



was used to stain the cell nuclei (blue).

### Immunohistochemistry

**Image 2.** Immunohistochemical analysis of Cytokeratin 17 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



### Western Blotting

**Image 3.** Western blot analysis of Cytokeratin 17 expression in PC12 (A), HeLa (B), A431 (C), NIH3T3 (D) whole cell lysates.