

Datasheet for ABIN7296184

anti-LIM Domain Kinase 1 antibody (Center)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	LIM Domain Kinase 1 (LIMK1)
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LIM Domain Kinase 1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human LIMK1.
Specificity:	Recognizes endogenous levels of LIMK1 protein.
Characteristics:	Rabbit polyclonal antibody to LIMK1
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	LIM Domain Kinase 1 (LIMK1)
Alternative Name:	LIMK1 (LIMK1 Products)
Background:	LIMK, LIM domain kinase 1, LIMK-1

Target Details

Gene ID:	3984, 16885
UniProt:	P53667 , P53668 , P53669
Pathways:	Caspase Cascade in Apoptosis , Regulation of Cell Size , CXCR4-mediated Signaling Events


Application Details

Application Notes:	WB (1:500 - 1:1000), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence

Image 1. Immunofluorescent analysis of LIMK1 staining in COLO205 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell



nuclei (blue).

Western Blotting

Image 2. Western blot analysis of LIMK1 expression in COLO205 (A), HEK293T (B), A431 (C), NIH3T3 (D), HeLa (E), HepG2 (F) whole cell lysates.