antibodies -online.com







anti-LIMK2 antibody (Center)





()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Overview	
Quantity:	100 μL
Target:	LIMK2
Binding Specificity:	Center
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LIMK2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human LIMK2.
Specificity:	Recognizes endogenous levels of LIMK2 protein.
Characteristics:	Rabbit polyclonal antibody to LIMK2
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	LIMK2
Alternative Name:	LIMK2 (LIMK2 Products)

Target Details

Background:	LIM domain kinase 2, LIMK-2	
Gene ID:	3985	
UniProt:	P53671	


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months


Images



Western Blotting

Image 1. Western blot analysis of LIMK2 expression in Jurkat (A), COLO205 (B), MCF7 (C) whole cell lysates.





Immunofluorescence

Image 2. Immunofluorescent analysis of LIMK2 staining in COLO205 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Immunohistochemistry

Image 3. Immunohistochemical analysis of LIMK2 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.