

Datasheet for ABIN7296392
anti-NDUFA8 antibody (Center)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	NDUFA8
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat, Cow, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NDUFA8 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NDUFA8.
Specificity:	Recognizes endogenous levels of NDUFA8 protein.
Characteristics:	Rabbit polyclonal antibody to NDUFA8
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	NDUFA8
Alternative Name:	NDUFA8 (NDUFA8 Products)
Background:	NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 8, Complex I-19kD, CI-19kD,

Target Details

Complex I-PGIV, CI-PGIV, NADH-ubiquinone oxidoreductase 19 kDa subunit

Gene ID: 4702, 68375

UniProt: [P51970](#), [Q9DCJ5](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of NDUFAB1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of NDUFA8 expression in HepG2 (A), MCF7 (B) whole cell lysates.