



[Go to Product page](#)

Datasheet for ABIN7296522  
**anti-PAWR antibody (C-Term)**

2 Images

Overview

Quantity:	100 µL
Target:	PAWR
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Dog, Chicken, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PAWR antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PAWR.
Specificity:	Recognizes endogenous levels of PAWR protein.
Characteristics:	Rabbit polyclonal antibody to PAWR
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	PAWR
Alternative Name:	PAWR ( <a href="#">PAWR Products</a> )
Background:	PAR4, PRKC apoptosis WT1 regulator protein, Prostate apoptosis response 4 protein, Par-4

## Target Details

Gene ID: 5074, 114774, 64513

UniProt: [Q96IZ0](#), [Q925B0](#), [Q62627](#)

## Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200)

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

## Images



### Immunohistochemistry

**Image 1.** Immunohistochemical analysis of PAWR staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



### Western Blotting

**Image 2.** Western blot analysis of PAWR expression in HeLa (A), NIH3T3 (B) whole cell lysates.