

Datasheet for ABIN7296704
anti-ERK1/2 antibody (C-Term)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	ERK1/2 (MAPK1/3)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Cow, Pig, Dog
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ERK1/2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human ERK1/2.
Specificity:	Recognizes endogenous levels of ERK1/2 protein.
Characteristics:	Rabbit polyclonal antibody to ERK1/2
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	ERK1/2 (MAPK1/3)
Alternative Name:	ERK1/2 (MAPK1/3 Products)
Background:	MAPK3, ERK1, PRKM3, Mitogen-activated protein kinase 3, MAP kinase 3, MAPK 3, ERT2,

Target Details

Extracellular signal-regulated kinase 1, ERK-1, Insulin-stimulated MAP2 kinase, MAP kinase isoform p44, p44-MAPK, Microtubule-associated protein 2 kinase, p44-ERK1, MAPK1, ERK2, PRKM1, PRKM2, Mitogen-activated protein kinase 1, MAP kinase 1, MAPK 1, ERT1, Extracellular signal-regulated kinase 2, ERK-2, MAP kinase isoform p42, p42-MAPK, Mitogen-activated protein kinase 2, MAP kinase 2, MAPK 2

Gene ID: 5595, 26417, 26413

UniProt: [P27361](#), [P28482](#), [Q63844](#), [P63085](#), [P21708](#), [P63086](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

Expiry Date: 12 months



Immunohistochemistry

Image 1. Immunohistochemical analysis of ERK1/2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Western Blotting

Image 2. Western blot analysis of ERK1/2 expression in HeLa (A), NIH3T3 (B), PC12 (C) whole cell lysates.

