antibodies - online.com







anti-RPL28 antibody (Center)





()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Overview		
Quantity:	100 μL	
Target:	RPL28	
Binding Specificity:	Center	
Reactivity:	Human, Monkey	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This RPL28 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)	
Product Details		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human RPL28.	
Specificity:	Recognizes endogenous levels of RPL28 protein.	
Characteristics:	Rabbit polyclonal antibody to RPL28	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Target Details		
Target:	RPL28	
Alternative Name:	RPL28 (RPL28 Products)	
Background:	60S ribosomal protein L28	

Target Details

Gene ID:	6158
UniProt:	P46779


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)	
Restrictions:	For Research Use only	

Handling


Format:	Liquid	
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.	
Expiry Date:	12 months	

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of RPL28 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of RPL28 expression in HeLa (A), MCF7 (B), HEK293 (C) whole cell lysates.