

Datasheet for ABIN7297186

anti-p53 antibody (C-Term, pSer315)



[Go to Product page](#)

3 Images

Overview

Quantity:	100 µL
Target:	p53 (TP53)
Binding Specificity:	C-Term, pSer315
Reactivity:	Human, Rat, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This p53 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human p53.
Specificity:	Recognizes endogenous levels of p53 (pS315) protein.
Characteristics:	Rabbit polyclonal antibody to p53 (pS315)
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	p53 (TP53)
Alternative Name:	p53 (TP53 Products)

Target Details


Background:	P53, Cellular tumor antigen p53, Antigen NY-CO-13, Phosphoprotein p53, Tumor suppressor p53
Gene ID:	7157, 24842
UniProt:	P04637 , P10361
Pathways:	p53 Signaling , MAPK Signaling , PI3K-Akt Signaling , Apoptosis , AMPK Signaling , Chromatin Binding , ER-Nucleus Signaling , Positive Regulation of Endopeptidase Activity , Hepatitis C , Protein targeting to Nucleus , Autophagy , Warburg Effect

Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500), IP (1:10 - 1:100)
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months




Immunofluorescence

Image 1. Immunofluorescent analysis of p53 (pS315) staining in A431 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



Immunohistochemistry

Image 2. Immunohistochemical analysis of p53 (pS315) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 3. Western blot analysis of p53 (pS315) expression in HeLa (A), A431 (B) whole cell lysates.