

Datasheet for ABIN7297314
anti-ZNF187 antibody (Center)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	ZNF187
Binding Specificity:	Center
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF187 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ZNF187.
Specificity:	Recognizes endogenous levels of ZNF187 protein.
Characteristics:	Rabbit polyclonal antibody to ZNF187
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	ZNF187
Alternative Name:	ZNF187 (ZNF187 Products)
Background:	ZNF187, Zinc finger and SCAN domain-containing protein 26, Protein SRE-ZBP, Zinc finger

Target Details

	protein 187
Gene ID:	7741, 432731
UniProt:	Q16670 , Q5RJ54


Application Details

Application Notes:	WB (1:500 - 1:1000), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence

Image 1. Immunofluorescent analysis of ZNF187 staining in HL60 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell



nuclei (blue).

Western Blotting

Image 2. Western blot analysis of ZNF187 expression in HL60 (A), SP2/0 (B) whole cell lysates.