# antibodies -online.com





Datasheet for ABIN7297398

## anti-IKBKG antibody (Center)

3 Images



Go to Product page

Overview	
Quantity:	100 μL
Target:	IKBKG
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat, Rabbit, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IKBKG antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human IKK gamma.
Specificity:	Recognizes endogenous levels of IKK gamma protein.
Characteristics:	Rabbit polyclonal antibody to IKK gamma
Purification:	The antibody was purified by immunogen affinity chromatography.
Target Details	
Target:	IKBKG
Alternative Name:	IKK gamma (IKBKG Products)

### **Target Details**

Background:	FIP3, NEMO, NF-kappa-B essential modulator, NEMO, FIP-3, IkB kinase-associated protein 1, IKKAP1, Inhibitor of nuclear factor kappa-B kinase subunit gamma, I-kappa-B kinase subunit gamma, IKK-gamma, IKKG, IkB kinase subunit gamma, NF-kappa-B essential modifier
Gene ID:	8517, 16151, 309295
UniProt:	Q9Y6K9, 088522, Q6TMG5
Pathways:	NF-kappaB Signaling, RTK Signaling, TCR Signaling, TLR Signaling, Fc-epsilon Receptor Signaling Pathway, Activation of Innate immune Response, M Phase, Production of Molecular Mediator of Immune Response, Hepatitis C, Protein targeting to Nucleus, Toll-Like Receptors Cascades, BCR Signaling, Ubiquitin Proteasome Pathway, S100 Proteins


## **Application Details**


Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months







#### **Immunofluorescence**

Image 1. Immunofluorescent analysis of IKK gamma staining in HL60 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

#### **Immunohistochemistry**

**Image 2.** Immunohistochemical analysis of IKK gamma staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **Western Blotting**

Image 3. Western blot analysis of IKK gamma expression in Jurkat (A), HL60 (B), HeLa (C), NIH3T3 (D), Raw264.7 (E), PC12 (F), rat heart (G) whole cell lysates.