antibodies -online.com





anti-CYP4F2 antibody (N-Term)

2 Images



Go to Product page

O	:
	view
	VICVV

Overview	
Quantity:	100 μL
Target:	CYP4F2
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CYP4F2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of
	human Cytochrome P450 4F2.
Specificity:	Recognizes endogenous levels of Cytochrome P450 4F2 protein.
Characteristics:	Rabbit polyclonal antibody to Cytochrome P450 4F2
Purification:	The antibody was purified by immunogen affinity chromatography.
Target Details	
Target:	CYP4F2
Alternative Name:	Cytochrome P450 4F2 (CYP4F2 Products)
Background:	Leukotriene-B(4) omega-hydroxylase 1, CYPIVF2, Cytochrome P450 4F2, Cytochrome P450-

Target Details

	LTB-omega, Leukotriene-B(4) 20-monooxygenase 1
Gene ID:	8529
UniProt:	P78329
Pathways:	Monocarboxylic Acid Catabolic Process


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of Cytochrome P450 4F2 staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pretreated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained



with haematoxylin and mounted with DPX.

Western Blotting

Image 2. Western blot analysis of Cytochrome P450 4F2 expression in A549 (A), HepG2 (B) whole cell lysates.